

УДК 547.29 612.017.34

## RESPIRATORY DYSFUNCTION OF PATIENTS WITH SEVERE BURN INJURY

*D.Kh Khaitov, I.I Toshev*

**Summary:** Dysfunctional changes in the upper respiratory tract in most cases develop as a post-burn pathology. (1,2,12). After a burn, in 30-40% of patients, the general condition worsens and vital organs fail. (3,6,8,11,13,16). The most important factors that damage the mucous membrane of the respiratory tract during IT are various combustion products in a confined space. The development of mucosal edema is primarily dependent on mediators such as histamine, bradykinin, prostacyclin and bronchospasm, and pulmonary vascular resistance is associated with the release of thromboxane and complement components (8.15).

**Key words:** Multiple organ disorders, burn disease, inhalation injury.

## ДЫХАТЕЛЬНАЯ ДИСФУНКЦИЯ БОЛЬНЫХ ТЯЖЕЛОЙ ОЖГОВОЙ ТРАВМОЙ

*Д.Х.Хайитов, И.И.Тошев*

**Резюме:** Дисфункциональные изменения верхних дыхательных путей в большинстве случаев развиваются как послеожоговая патология. (1,2,12). После ожога у 30-40% больных ухудшается общее состояние и отказывают жизненно важные органы. (3,6,8,11,13,16). Важнейшими факторами, повреждающими слизистую оболочку дыхательных путей при ИТ, являются различные продукты горения в замкнутом пространстве. Развитие отека слизистой оболочки зависит в первую очередь от медиаторов, таких как гистамин, брадикинин, простаглицин и бронхоспазм, а легочное сосудистое сопротивление связано с высвобождением тромбосана и компонентов комплемента (8.15).

**Ключевые слова:** Полиорганных нарушения, Ожоговой болезни, ингаляционная травма.

## YUQORI NAFAS YO`LLARINING KUYISHDAN KEYINGI FUNKSIONAL O`ZGARISHLARI

*D.H Hayitov, I.I Toshev*

**Резюме:** Yuqori nafas yo`llarining disfunktsional o`zgarishlari ko`pchilik holatlarda kuyishdan keyingi patologiya sifatida rivojlanadi. (1,2,12). Kuyishdan keyin 30-40% bemorlar umumiy ahvoli og`irlashib ulardagi hayotiy muhim a`zolar faoliyati izdan chiqishi kuzatiladi. (3,6,8,11,13,16). IT vaqtida nafas yo'llarining

shilliq qavatiga zarar etkazadigan eng muhim omillar - bu yopiq makonda turli xil yonish mahsulotlari. Rivojlanayotgan shilliq qavat shishi, birinchi navbatda, gistamin, bradikinin, prostatsiklin va bronxospazm kabi vositachilarga bog'liq va o'pka tomirlarining qarshiligi tromboksan va komplement komponentlarining chiqishi bilan bog'liq (8.15).

**Kalit soʻzlar:** Poliorgan yetishmovchilik, kuyish kasalligi, ingalyatsion jarohat.

Acute respiratory failure in burned patients, regardless of the cause that caused it, often leads to multiple organ failure and mortality. Respiratory failure is one of the most common forms of organ failure in the PON syndrome and, progressing, leads to death in combination with dysfunction of other organs, regardless of the presence of inhalation injury (12,16). The disorder of the respiratory function of the lungs, in turn, disrupts the nutrition of the alveolar tissue, reduces the production of surfactant, contributing to the development of atelectasis (4.14), and increases venous blood shunting (2.5). In burn disease, along with respiratory disorders, non-respiratory functions of the lungs suffer: blood purification from mechanical impurities and their enzymatic destruction, participation in hemostasis and fibrinolysis, water and electrolyte metabolism, metabolism of proteins, fats, biologically active substances (1, 7,9,10).

**Purpose of the study:** To study the pathology of the respiratory organs in burn disease

**Materials and methods of research:** For the period 2017-2021, there were 2136 patients in the combustiology department of the Bukhara branch of the RRCEM. Of these, 156 patients had impaired function of the respiratory system. Of these, 96 women (61.5), men 60 (38.5). The age of the patients ranged from 2 to 70 years.

Frank index (IF) more than 60-90 units in 76 (48.7) patients, IF more than 90 units in 80 (51.3%) patients. In 55% burns with flames, 43% with boiling water, in 2% - electrical injury. ) To study the dysfunction of the respiratory system in terms of systemic inflammatory (SIR), we analyzed the clinical indicator of tachypnea and features of the X-ray picture of the lungs (Table No. 1).

Chest radiography is an important diagnostic method for monitoring pulmonary complications and evaluating treatment from day 1 of a burn injury (493).

**Results and discussions:** First of all, it was noted that among 156 patients with identified disorders of the respiratory system, the vast majority (128 patients) with tachypnea 21-24 in 1 minute had signs of venous plethora of the lungs, expressed to varying degrees and indicating an increase in pressure in the pulmonary artery system. According to the literature data, the pressure in the pulmonary artery is increased during all periods of burn disease, depending on the severity of the burn disease, and the greatest increase in pressure was found in the stages of shock and toxemia with a gradual decrease by the time of healing of burn wounds (11.16).

Hyperserotopinemia plays a role in the genesis of increased pulmonary resistance, hypertension of the pulmonary circulation, impaired perfusion and gas exchange, causing vasoconstriction and bronchial constriction (2,4). The development of venous plethora, and with it tachypnea in burned patients, we associate with manifestations of systemic microcirculation disorders and capillary-alveolar diffusion of blood gases against the background of developed SIRS (17.18.19.20).

Table №.1

Pathological changes	Number of patients в Г]			groups	Total 156
	1-18	II-40	III - 56	IV-42	
Venous plethora	7	30	52	39	128
Interstitial edema	-	3	10	10	23
Alveolar edema	-	1	3	6	10
ARDSV	-	-	1	2	3
bacterial pneumonia	-	1	8	11	20
Infarct pneumonia	1	5	6	3	15
Atelectasis	1	3	8	6	18
Diaphragm dome position	-	5	7	8	20
Pneumothorax	-	-	1	-	1

Distribution of patients with different severity of burn injury in accordance with x-ray changes in the lungs in most patients, signs of venous plethora appeared on the 1-2 day and persisted for 7-14 days, and in some patients it was much longer during the SVR, which correlated with the severity of the burn injury. With an uncomplicated course of a burn disease, venous plethora, according to radiography, gradually decreased and disappeared, with a complicated one, it appeared again or increased, while some patients developed signs of increased vascular permeability in the microvessels of the lungs, which manifested itself as interstitial or even alveolar edema. Among 128 patients, more than half (5, 8, 11) had tachypnea combined only with circulatory disorders in the lungs.

Signs of increased vascular permeability against the background of venous plethora in the form of interstitial and alveolar pulmonary edema were detected in 23 and 10 patients, respectively. An additional negative role in the formation of edema can be played by excessive infusion therapy.

**Conclusions:** We assume that interstitial edema, caused by a systemic increase in vascular permeability in SVR, occurred in a larger number of patients with tachypnea, but its severity was minimal and insufficient to detect this sign on radiographs. A more

pronounced swelling of the interstitium, and even more so the sweating of edematous fluid into the alveoli, caused more severe dysfunction and even respiratory failure.

### Literature:

1. Барвинский, В.Н. Влияние ожоговой травмы на функциональное состояние сердечно-сосудистой системы в процессе хирургического лечения: Дис. ... канд. мед. наук.- Киев, 1989.- 144 с.
2. Бугров, С.Н. Нарушение серотонин-регулирующей функции легких в остром периоде экспериментальной ожоговой болезни / С.Н. Бугров // Травматология и ортопедия России.- 1996.- N 1,- С.40-41.
- 3.Вазина, И.Р. Функциональная морфология легких при термической травме: Дис. ...д-ра мед. наук.- М., 1988.-360 с.
- 4.Дороватовский, В.А. Состояние сурфактантной системы легких при острой дыхательной недостаточности / В.А. Дороватовский, А.Х. Мустафин // Анестезиология и реаниматология.- 1986.- N 4.- С.23-25.
- 5.Зильбер, А.П. Дыхательная недостаточность. Руководство для врачей. /П.Зильбер.- М.:Медицина, 1989.- 512 с.
6. Калашников, А.Ю. Эндоскопическая диагностика поражений слизистой оболочки верхнего отдела желудочно-кишечного тракта у больных с термическими ожогами тела: Автореф. дис. ...канд. мед. наук. - М., 2005. - 22 с.
- 7.Клевцова, Т.М. Функциональное состояние сердечно-сосудистой системы у обожженных: Дис... канд. мед. наук. - Ленинград, 1988.-170 с.
8. Очилова Д.А., Комилова Б.О. Распространенность факторов риска сердечно-сосудистых заболеваний среди населения//Кардиология Узбекистана.-2018.-№2(48).-С.21-23.
- 9.Логинов, Л.П. Трахеостомии у больных с тяжелой ингаляционной травмой / Л.П.Логинов // Проблемы лечения тяжелой термической травмы: мат. VIII Всерос. Пауч.-практ. конф. с междунар. участием,- Н.Новгород, 2004.- С.86-87.
- 10.Сыромятникова, Н.В. Метаболическая активность легких / Н.В. Сыромятникова, А.В.Гончарова, Т.В.Котенко.- Л.: Медицина, 1987.- 165 с.
11. Khayrullaeva D. Kh., Avezova S.M. Changes in hormonal in adverse environmental condition//World Journal of Pharmaceutical Research 2020 Vol.9 Issue 5, pp. 2450-2458.
12. Хайруллаева Д.Х. Состояние дисциркуляторной энцефалопатии при гипотериозе// Tibbiyotda Yangi kun O`zbekiston.-2020.-№.2/1(30/1).-С. 69-71.
13. Inhalation injury in burned patients: effects and treatment / D.N. Herndon, R.E. Barrow, H.A. Linares et al. // Burns. - 1988. - N 14.- 349-356.
14. Очилова Д.А., Рўзиев О.А., Комилова Б.О. Юрак қон томир касалликлари ривожланишида носоглом турмуш тарзининг аҳамияти//Ўзбекистон Кардиологияси.-2018.-№2(48).-С.34-35.

15. Schulz, A.-M. Pathophysiologie, Diagnose und Therapie des Inhalatio-  
straumas / A.-M. Schulz, A. Werba // Acta Chir. Austriaca.-1997.-Vol.29.-N 6,- P.312-  
315.

16. role of inhalation injury in bum trauma / E.E. Tredget, H.A.Shankowsky, T.V.  
Taerum et al. // Ann. Surg.-1990.-Vol. 212.- N 6.- P. 720727.

17. Usmonov U.R, SobirovSh,H, Irgashev I,E. The role of  
neuroimmunoendocrine processes in the human body// World Journal of  
Pharmaceutical Research Belgium-2020.-Volume9.-Issue. 62413-2424 pages.

18.USMONOV U.R, IRGASHEV I,E, CHANGES IN THE  
MORPHOFUNCTIONAL PROPERTIES OF THYMUS AND SPLEEN UNDER THE  
INFLUENCE OF MITES OF DIFFERENT ORIGINS”//Тиббиётда янги кун  
Ўзбекистан.-2020.-№2(30).-С. 242-245.

19. Очилова Д.А., Рузиев О.А., Худойберганова Н.Х., Бобоев А.Т. Оценка  
эффективности антигипертензивной препаратов//Вестник медицинского центра  
управления делами президента республики Казахстан. - 2015. - №3(60). – С. 41-  
44.

20. Ochilova D.A., Komilova B.O. Value of the pyridoxine of the hydrochloride  
and copper in prevention of atherosclerosis(Literary review)// East European Scientific  
Journal. Warszawa. Polska. 2019. – P. XX-XX.