

УДК 547.29 612.017.34

**SPECIFICITY OF RESUSCITATION MEASURES IN PATIENTS WITH ISCHEMIC HEART DISEASE AND ARRHYTHMIA**

**Khayitov Dilshod Khayotovich\***; **Toshev Izzat Ikromovich**

Assistant, Department of Surgical Diseases and Resuscitation,  
Bukhara State Medical Institute named after Abu Ali ibn Sina,

**Abstract**

The situation after cardiopulmonary resuscitation (CPR) is one of the most complex and unresolved medical problems, with world statistics showing that more than 61% of patients who undergo CPR die within the first 22 hours. Although 40% of patients survived, various complications were observed, of which only 16% of patients fully recovered, and the remaining 84% of patients retained various complications [1-8]. Despite the improvement of the pulmonary-cardiac-cerebral resuscitation (PCCR) algorithm, the lack of knowledge and practical experience in conducting PCCR to patients has led to a high mortality rate of patients and a high level of complications. In some sources it is known that in cases of late onset of pulmonary heart resuscitation, the patient's vital functions are fully restored, depending on the patient's age, ambient temperature, resistance to hypoxia, first aid provided by non-medical personnel around the patient [9-18].

**Objective of the study:** To study modern intensive care in the treatment of post-resuscitation complications resulting from cardiac arrhythmias.

**Keywords:** Post-Resuscitation Disease, Cardiac Arrhythmias, Cardiopulmonary Resuscitation, Hypoxia, Oxygen Therapy, Neuroprotection.

**INTRODUCTION**

**Case study:** Patient A., 30 years old, was brought to the Bukhara branch of the Republican Scientific Center of Emergency Medicine (BBRSCEM) on December 26, 2021 at 23:10 in a road car in case of clinical death.

From the anamnesis, the patient has been drinking alcohol and energy drinks regularly during the last 4-5 days while attending wedding ceremonies. 20 minutes before arriving at the hospital, he

came out of the kitchen and collapsed and fainted while talking to his colleague. The patient was transported by colleagues to the BBRSCEM in his car on 26.12.2021 at 23:10 and was taken to the shock ward of the admission department. Pulmonary cardiopulmonary resuscitation was started immediately due to the absence of pulsation in the central carotid arteries in the shock chamber, shortness of breath, asystole in the cardiomonitor. The patient's heart function was restored in 20 minutes. The patient's hemodynamics was transferred to the Department of Surgical Resuscitation at 01:30

on 27.12.2021 in case of maintenance under the influence of high-dose vasopressor drugs.

From the anamnesis of life, the patient denies hereditary diseases, which he grew up in satisfactory conditions. Does not consume regular medications.

When he arrives, the patient is not conscious. Body structure is normostenic, subcutaneous fat layer is moderately developed. The skin and visible mucous membranes are cyanotic. There are no scars on the body. Deformations in the musculoskeletal system are not detected. Temperature 36.0 OS. Maximum pupil dilation OD = OS, photoreaction is not called. Atonia. Areflexia. Breathing-abs. Saturation - abs. Heart sounds are not audible, no pulse is detected in the central arteries. Asystole is detected in the cardiomonitor. ABP-not detected. The tongue is wet, clean. There are no vomit masses in the oral cavity. The abdomen is soft, the liver and spleen cannot be felt. Involuntary defecation and urinary incontinence are not detected.

In the shock chamber of the reception department of the BBRSCEM reception of the patient, there was no pulsation in the central carotid arteries, no breathing, asystole in the cardiomonitor. External cardiac massage was initiated and tracheal intubation was performed using an bu7.5 intubation tube while simultaneously breathing with an Ambu copy. Artificial Lung Ventilation(ALV) apparatus Hamelton C3, A / S Volume mode. DO-600 ml, f-18 times per minute, PSV-14 mambr, PEEP-2, FiO<sub>2</sub> -100%. In order to restore cardiac function, r-r Adrenaline-0.18% -1.0, r-r Atropine sulfate-0.1% -1.0 were used 3 times in 5 minutes. For maintenance of blood pressure and vasopressor therapy, r-r Dexamethasone-8 mg every 5 minutes, r-r Dopamine 40 mkg.kg.min., R-r Noradrenaline-2,0 mkg.kg.min. started at. Due to the observed ventricular fibrillation on the cardiomonitor at 18-19 minutes of pulmonary-cardiac resuscitation, 3 defibrillations + external cardiac massage + defibrillation + external cardiac massage + defibrillation were performed with a current of 200-250-300 Dj. At 20 minutes, after 3 defibrillations, normal heart function was restored. The patient's heart function was restored in 20 minutes. As the patient's vital signs began to breathe with resistance to the ALV apparatus, sodium oxybutyrate-4 g, Sibazon-10 mg, morphine hydrochloride-10 mg were administered intravenously, the patient was adapted to the ALV apparatus and transferred to the intensive care unit against vasopressors.

On arrival Total blood test: hemoglobin-123 g / l; erythrocytes-4.0 x 10<sup>12</sup> / l; color index-0.92; hematocrit-37%; blood clotting time: start-2:58, end-3:24; platelet-133.0 x 10<sup>9</sup> / l; leukocyte-5.3 x 10<sup>9</sup> / l; rod nucleus neutrophils-2%; segment nuclear neutrophils-38%; Eosinophils-2%; Lymphocytes-51%; monocytes-7%; erythrocyte sedimentation rate-4 mm / s. Biochemical blood test: Total protein-58 g / l; glucose-7.1 mmol / l; urea-6.0 mmol / l; bilirubin: total-29.0 mmol / l, bound-9.0 mmol / l; ALT-277 Ed / l; AST-291 Ed / l; Calcium-2.3 mmol / l; fibrinogen-4.4 g / l. General urine analysis: color-yellow, clear; relative density-1020; rN-7.0; protein-0.066 g / l;

glucose-negative; ketone bodies-positive (+); blood unit detection reaction-positive; bilirubin-negative; epithelial cells in a drying area - flat-0-1; leukocytes -8-9 in a drying area; erythrocytes unchanged in a drying area-6-7. Chest X-ray findings: Bronchitis; Conclusion of abdominal ultrasound examination: Parenchymatous organ injury and free fluid in the abdomen were not detected at the time of examination; Conclusion of cerebral MSKT examination: No organic or traumatic structural changes in brain tissue were detected.

### TABLE 1 HEMOGLOBIN AND HEMATOCRIT CONDITION

As a result of intensive therapy, clinical and laboratory parameters improved in dynamics: In the analysis of whole blood: hemoglobin-121 g / l; erythrocytes-4.1 x 10<sup>12</sup> / l; color index-0.9; hematocrit-39%; coagulation time: start-4:10, end-4: 30; platelet-176.0 x 10<sup>9</sup> / l; leukocyte-8.5 x 10<sup>9</sup> / l; rod nucleus neutrophils-2%; segment nuclear neutrophils-70%; Eosinophils-1%; Lymphocytes-25%; monocytes-2%; erythrocyte sedimentation rate-8 mm / s. Biochemical blood test: Total protein-70 g / l; glucose-6.4 mmol / l; urea-6.0 mmol / l; bilirubin: total-11.4 mmol / l, bound-1.8 mmol / l; ALT-95 Ed / l; AST-57 Ed / l; Calcium-2.2 mmol / l; fibrinogen-3.8 g / l. General urine analysis: color-yellow, clear; relative density-1016; protein traces; glucoseneegative; ketone bodies-negative (-); blood unit detection reaction-negative; bilirubin - negative; epithelial cells in a drying area - flat-0-1; leukocytes -4-6 in a drying area; erythrocytes unchanged in a drying area-1-2. Chest X-ray findings: Bronchitis; Conclusion of abdominal ultrasound examination: No echopathology was detected in the abdomen at the time of examination; Conclusion of cerebral MSKT examination: No organic or traumatic structural changes in brain tissue were detected. ECG: Sinus rhythm. YuQS-84 times per minute. Mild metabolic changes were detected in the myocardium.

The patient continued intensive care in the surgical intensive care unit from 01:30 on 27.12.2021. The patient's hemodynamics were captured under the influence of vasopressors. The patient was discontinued vasopressor therapy at 14:00 on 30.12.2021 against the background of a decrease in the dose of vasopressors. The patient's consciousness was completely restored from coma II to 29.12.2021 to coma I, 31.12.2021 to sopor level, 02.01.2022 to mild numbness, 03.01.2022. The patient's consciousness was maintained in constant analgo-sedation until it improved to a mild level of numbness and was awakened from time to time to determine the level of consciousness. The patient underwent tracheostomic tube implant surgery on 29.12.2022 in order to improve the sonication of the tracheobroncheal tree, reduce the volume of the dead cavity, to prevent disruption of the oral microflora. The patient was transferred to independent breathing on 03.01.2022 through a tracheostomy. On 05.01.2022, the tracheostomy tube was removed and the ligament was burned. The patient was voluntarily discharged home on 06.01.2022 without cognitive dysfunction.

Based on the results of objective examination and laboratory-instrumental examination, the following diagnosis was made: Acute toxic myocarditis of alcoholic etiology? Complications: Ventricular fibrillation. Asystole. Condition after pulmonary heart-brain resuscitation. Postreanimation disease. Keeping the patient in complete hypothermia to the treatment plan; for neuroprotection: citicoline; In order to improve metabolism in organs and tissues: cytoflavin; In order to maintain and improve the activity of brain cells remaining in ischemia: amantadine sulfate; for membrane stabilization: lidocaine, dexamethasone; for dehydration: hypertonic sodium chloride solution, furasemide; For antioxidant therapy: Ascorbic acid; For antibacterial therapy: ceftriaxone, levofloxacin; In order to improve the rheology of the mine: reosorbilact, heparin; For construction and power supply: Melkam; For nebulizer therapy: decasan, nebutamol; For analgo-sedation: Sodium oxybutyrate, profol, verzepam, midozalam, metamizole sodium, fentanyl, promedol, morphine hydrochloride; For muscle relaxation: Arduan; For the treatment of hyperthermia: Taken paracetamol.

**TABLE 3 PHARMACOTHERAPY IN POST-RESUSCITATION DISEASE**

The name of the drug	Day	December 2021	January 2022	26	27	28	29	30	31	1	2	3	4	5	6
Cyticolin															
Cytoflavin															
Amantadine sulfate															
Lidocaine															
Dexamethasone															
Sodium 7.5%															
-200.0 Furasemide															
Ceftriaxone															
Levofloxacin															
Reosorbilakt															
Heparin															
Nebulizer therapy															

**CONCLUSIONS:** 1. All over the world, it is accepted that the time norm of clinical death is 3-8 minutes, but due to the achievements of modern medicine, this time can be extended to 10-15 minutes. This could allow thousands of patients in the community to return to social life without cognitive dysfunction, without social death. Intensive therapies performed in 4 patients in our practice in patients who remained in a state of clinical death for 10-15 minutes gave a positive result.

2. Intensive therapy of post-resuscitation disease should be carried out in combination with treatment of post-anoxic encephalopathy and treatment of polyorgan deficiency, which develops in complex forms of post-resuscitation disease. 3. Treatment of postanoxic encephalopathy includes neurotropic therapy: a combination of antihypoxants, antioxidants, sedative drugs, and energy sources. 4. Restoration of brain activity after clinical death depends on maintaining cerebral perfusion in the 1-2 stages of the disease after resuscitation in a slightly higher than normal and early onset of neuroprotective drugs. 5. In patients with pulmonary resuscitation and the restoration of cardiac function, the initiation of cerebral therapy from the first minutes restores the function of nerve cells in hypoxia and ischemia, as well as a sharp decrease in the incidence of cognitive impairment.

**REFERENCES:**

1. Kuzovlev A.N. (2017). Extended resuscitation in circulatory arrest: current recommendations and prospective studies. Bulletin of Intensive Care named after A. I. Saltanov, (3), 53-57.
2. USMONOV U.R, IRGASHEV I.E, CHANGES IN THE MORPHOFUNCTIONAL PROPERTIES OF THYMUS AND SPLEEN UNDER THE INFLUENCE OF MITES OF DIFFERENT ORIGINS”//Тиббиётда янги кун Ўзбекистан.-2020.-№2(30).-С. 242-245.
3. Vismont, F. I. Pathological physiology: textbook / F. I. Vismont [and others]; under the editorship of prof. F.I. Vismont. - 2nd ed., revised. - Minsk: Higher School, 2019 - 640 S. : ill.
4. Wismont, F.I. General pathophysiology: textbook / F.I. Vismont, E.V. Leonova, A. V. Chanturia. - Minsk: Higher School., 2011 - 364 p.
5. Evtushenko, S. V. Analysis of the effectiveness of resuscitation / S. V. Evtushenko, S. S. Grachev, N. K. Ivankovich // Medical Journal. - 2015 - No. 3 - S. 61-65.
6. Usmonov Ulugbek Rizoyevich. (2022). Morphological and Functional Aspects of the Lymphatic System and its Current Problems. Eurasian Journal of Research, Development and Innovation, 11, 5–8. Retrieved from <https://www.geniusjournals.org/index.php/ejr/article/view/2054>
7. Usmonov U.R. (2022). Cognitive Dysfunction After Multiple Surgery. Eurasian Medical Research Periodical, 9. Retrieved from <https://www.geniusjournals.org/index.php/emrp/article/view/1678>
8. Usmonov Ulug`bek Rizoyevich, Jakhonov Obidjon Olimjonovich, Sobirov Shokhrukh Khusenovich, Khaitova Dildora Sharifboevna, CHANGES IN THE MORPHOFUNCTIONAL PROPERTIES OF THYMUS, SPLEEN AND LYMPHOID SYSTEM UNDER THE INFLUENCE OF MITES OF DIFFERENT ORIGINS , Web of Scientist: International Scientific Research Journal: Vol. 3 No. 1 (2022): was
9. Rizaeva, M. Zh. (2020). Efficiency And Safety Of Electric Cardioversion In Persistent Form Of Atrial Fibrillation. A new day in medicine, (4), 322-325.
10. Kenzhaev, M. L., & Rizaeva, M. Zh. (2020). Clinical Course Of Atrial Fibrillation In Patients With Coronary Heart Disease. In Science and Innovation-Modern Concepts (pp. 103-109).
11. Turdiev, U. M., Boltaev, E. B., & Kodirov, M. D. (2020). Cytokine Indicators In Patients With Acute Coronary Syndrome Depending On The Type Of Antithrombotic Therapy. In Graduate School: Research (pp. 93-97).
12. Urakov, Sh. T., Eshonov, O. Sh., & Boltaev, E. B. (2019). Postoperative Cognitive Dysfunction. A new day in medicine, (2), 56-60.

- 13.** Eshonov, O. Sh., & Amonov, S. Zh. (2021). Hypocalcemia as a Factor In The Development Of Cognitive Dysfunction After Thyroid Surgery. *Scientific progress*, 1(5).
- 14.** Eshonov, O. Sh., & Kayumov, M. (2021). Features of cognitive dysfunction after multiple operations. *Scientificprogress*, 1(5).
- 15.** Eshonov, O. Sh., Oltiev, U. B., & Zhamolov, M. M. (2018). Postoperative cognitive disunction. *National Journal Of Neurology*, 39-41. **16.** Sabirov, D. M., Eshonov, O. Sh., Batirov, U. B., & Khaidarova, S. E. (2017). Postoperative cognitive dysfunction. *Bulletin of emergency medicine*, (2).
- 17.** Kenzhaev, M. L., & Rizaeva, M. Zh. (2020). Detection Of Predictors Of Atrial Fibrillation In Patients With Coronary Heart Disease. *A new day in medicine*, (2), 403-406.
- 18.** Koyirov, A. K., Shirinov, D. K., Rakhimov, M. M., Boltayev, E. B., & Galilov, A. A. (2021). Non-Invasive Lung Ventilation In Acute Respiratory Failure Caused By New Coronavirus Infection Covid-19. *Новый день в медицине*, (1), 107-114.
- 19.** Boltayev, E. B. (2021). Choice Of Respiratory Therapy In Severe Patients With New Coronavirus Infection Covid-19. *Достижения науки и образования*, (8 (80)), 70-74.
- 20.** Boltayev Elmurod Bekmurodugli, & Sabirov Joraboy Marifbayevich. (2021). Organization Of Oxygen Therapy Using A Nasal Mask And Ventura Mask In Severe Patients With Covid-19 Corona Virus Infection. *Emergent: Journal of Educational Discoveries and Lifelong Learning (EJEDL)*, 2(08), 6–10. <https://doi.org/10.17605/OSF.IO/N26VB>
- 21.** Ярашев, А. Р., Болтаев, Э. Б., & Шаббаев, Ж. К. (2020). A Retrospective Analysis Of Complications Of Percutaneous Dilatation Tracheostomy. *Новый день в медицине*, (4), 301-304.
- 22.** Rizaeva, M. Z. (2022). The Clinical Course Of Atrial Fibrillation In Patients With Coronary Heart Disease. *European journal of molecular medicine*, 2(1).