

MODERN TRENDS IN TREATMENT SICK MICROSPORIA

***Pak Uliana Ilyaevna***

*Cosmetologist of the Samarkand Regional Branch of the  
Republican Specialized Dermatovenerological and  
Cosmetology Scientific and Practical Medical Center*

***Tillakobilov Ismoil Bahodirovich***

*Assistant of the Department of Skin and Venereal Diseases,  
Samarkand State Medical University*

***Abdullaev Xasan Davlatovich***

*Assistant of the Department of Skin and Venereal Diseases,  
Samarkand State Medical University*

***Tolibov Mansur Mahmudovich***

*Assistant of the Department of Skin and Venereal Diseases,  
Samarkand State Medical University*

**Annotation:** Microsporia - This common infectious dermatomycosis, affecting the skin and its appendages in humans, as well as the skin and hair of animals. If 20-30 years ago, microsporia was recorded almost exclusively in children under 14 years, in the last 10 years there have been many publications on the detection of the disease in family foci with lesions of smooth skin, vellus and long hair in adults, including elderly of people [3, 5]. More often became register infiltrative suppurative microsporia pilaris parts heads [4].

**Keywords:** microsporia, treatment

**Introductions:** In Republic Uzbekistan, To unfortunately traditionally, more co times THE USSR, a relatively high incidence of scabies is revealed (more than 35-48 cases per 100thousand people). And in the last 5 years (2019-2023 ) , the incidence of microsporia in Republic Uzbekistan remained on relatively high level: V 2019 G - 41.3 cases / 100 thousand population, in 2020 - 45.0, in 2021 - 48.0, in 2022 - 40.5 and in 2023 38.6 cases / 100 thousand population. Experience of mycological laboratories Uzbekistanshows that currently in patients with microsporia of the scalp And smooth skin sown only M. lanosum (M. canis ). So, V mycological laboratory was received 923 culture M. lanosum at survey 771 a patient with microsporia (including all patients upon admission). We agree with V.M. Rukavishnikova (2001) that the fluffy microsporum has become a kind of cosmopolitan mushroom, practically the only pathogen zoophilic microsporia on Eurasian continent.

drug choice For treatment microsporia many years was griseofulvin (in many developed countries it still remains so). However, in the last 10 years in the treatment of patients with microsporia began to widely use new systemic antimycotics - terbinafine and itraconazole [1-5]. Because in the Republic Uzbekistan in the last 3

years, griseofulvin practically does not enter the pharmacy network we were also forced to switch to the treatment of patients microsporia preparations terbinafine ( itraconazole preparations for children under 12 years of age, the Pharmacopoeia of the Republic of Belarus does not recommend apply).

We compared the incidence of microsporia in different months of the year. IN period With 2019 G. By 2023 G. amplitude monthly hesitation incidence microsporia were very similar That's why we considered possible summarize by months these results for 5 years.

Table 1. The number of cases of microsporia in 2019-2023 at grouping by months

Jan- var	Feb - ral	March	Ap- relay	May	June	July	Av - thick	Saint- October	OK- October	But- Novem ber	De- Decem ber
162	140	61	64	70	60	70	204	556	490	294	248
363			194			830			1032		

Least incidence microsporia V recent 5 years was registered in March-July, and in August-September there was a pronounced increase registration cases microsporia With subsequent gradual decline this indicator With October By February. Observed seasonality epidemic process is due to the seasonality of the microsporia epizootic among cats (the main source of infection), as well as seasonal migration of the population, when in August-September children are returning V city And look around parents And health workers at their admission to schools and kindergartens. Sources of infection of patients with microsporia were detected in 98.0-99.5% of cases. Kittens predominated among the sources of infection. (more than 91.5% on average), cases of infection with zoophilic microsporia from others sick (up to 4.7 % V separate years).

**aim** present research is conducting comparative estimates clinical efficiency And economic costs at use methodologies treatment microsporia smooth skin, hairy parts heads And combined mycotic defeat hairy parts heads And smooth skin griseofulvin And drugs terbinafine ( lamisil , lamiter , exifin , terbizil ).

**MATERIALS AND METHODS**

In total, 272 patients with microsporia aged 2 years and over were under observation. up to 44 years, of which 144 (52.9%) are male and 128 (47.1%) are female. Aged before 4 years was 14.3 % sick, 5-7 years - 22.8 %, 8-14 years - 51.7 %, 15 - 18 years - 5.5 %, older 18 years - 5.7 %.

Microsporia of the smooth skin of the trunk and extremities was diagnosed in 70 patients, including 8 adults. Microsporia was diagnosed in 117 patients scalp, and in 85 - a combined lesion with fluffy microsporum hairy parts heads And smooth skin, V volume including at 3 patients including 1 adult patient, there was infiltrative

suppurative form of microsporia At all sick before treatment clinical diagnosis was confirmed laboratory (microscopically in 100%, mushroom culture microsporum lanosum was obtained from 98.7 % patients).

The number of foci on the scalp was as follows: 1 foci in 51.8%, 2 foci - in 30.1%, 3-4 foci - in 8.2%, 5 or more foci - in 9.9% of patients; on smooth skin, respectively, up to 3 foci were in 23.5%, 4-5 foci - in 12.8%, 6-10 foci - in 15.7 %, up to 20 foci - at 28.3%, over 20 foci - at 19.7 %.

Griseofulvin appointed By classical three-stage scheme. First 2-4week (until the first negative fungal test) the drug was administered daily from calculation of 22 mg / kg of body weight per day (daily dose was taken in three divided doses ). After the first negative analysis, the drug in the same dose was administered every other day, and after three consecutive negative tests (tests were performed 1 time in 5-7 days) griseofulvin V complete daily dose appointed 2 times V week more 2 weeks.

Terbinafine preparations were prescribed according to the classical method (it is recommended specialists from firms Novartis , \_ developed branded a drug terbinafine - lamisil ) at the rate of : children weighing up to 20 kg received 1 time per day 62.5 mg, with a body weight of 20 to 40 kg - 125 mg, over 40 kg - 250 mg; patients with massive lesions of the scalp and smooth skin received drugs terbinafine according to the method of Professor N.N. Potekaev from the calculation : children up to 20 kg - 94 mg, from 20 to 40 kg - 187 mg, and over 40 kg - 250 mg.

Therapy systemic antimycotics was carried out on background generally accepted external therapy (shaving and shampooing 2 times a week, 2-5% iodine tincture and fungicidal ointments). If within 3.5-5 weeks of treatment with terbinafine the results microscopic research on mushrooms were stable positive majority children translated With classical methods treatment on methodology treatment professors N.N.Potekaeva With 1.5 times increase daily doses terbinafine [ 5 ].

The criteria for the effectiveness of treatment were the rate of clinical resolution. foci defeat, terms registration first And third contract negative results microscopic research hemp broken off hair V outbreaks defeat.

## RESULTS AND THEIR DISCUSSION

For the treatment of patients with microsporia of smooth skin along with external therapy appointed systemic antimycotics : 6 sick - lamisil , 27- exifin , 37 - lamiter . On background treatment lamisil first negative analysis on Availability mycelium of the fungus in skin flakes and vellus hair was obtained at  $13.6 \pm 2.1$  days, and the third negative analysis - at  $23.2 \pm 2.4$  days. In patients treated with exifin , these figures were  $14.7 \pm 2.0$  and  $24.4 \pm 2.1$  days, respectively , and those who received lamiter -  $15.7 \pm 2.3$  and  $25.8 \pm 2.7$  days. Thus, the treatment of patients with microsporia smooth skin

With using tested systemic antimycotics terbinafine lasts an average of 3.5 weeks, results are somewhat preferable at use lamisila , on differences statistically Not reliable (P>0.1-0.5).

The results of microscopic examination of hair stumps for fungi in the process treatment of patients with microsporia of the scalp and combined lesions scalp and smooth skin of the trunk and extremities are reflected in the table 2.

Table 2. Dynamics mycological cure microsporia hairy parts heads at use various systemic antimycotics

Systemic antimycotic	Negative results of microscopic examination mushrooms hemp hair V process treatment (V days)					
	microsporia parts heads			combined defeat h/h heads And smooth skin		
	n	first negative analysis	third negative analysis	n	first negative analysis	third negative analysis
Griseofulvin	60	24.7+ _ 4.4	35.0+ _ 3.5	47	25.1+ _ 4.8	34.4+ _ 5.9
Lamisil	10	22.0+ _ 3.2	32.2+ _ 4.0	8	19.1+ _ 2.6	32.3+ _ 3.6
Lamiter	20	23.8+ _ 5.7	34.4+ _ 6.6	12	27.1+ _ 4.9	38.3+ _ 5.2
Exifin	16	26.1+ _ 3.5	37.3+ _ 3.6	9	26.7+ _ 5.0	36.2+ _ 6.1
Terbizil	elevation	26.8+ _ 4.5	38.2+ _ 5.1	9	27.5+ _ 4.3	40.1+ _ 5.3

As can be seen from the table, the terms of negative microscopic analyzes for fungi in stumps broken off hair at microscopy hairy parts heads And at combined damage to the scalp and smooth skin when using griseofulvina And drugs terbinafine comparable differences statistically unreliable. Comparison of the terms of microbiological treatment of microsporia with damage to the scalp, as well as the scalp and smooth skin torso And limbs various drugs terbinafine showed What branded a drug lamisil accelerated microbiological cure V average on 2-8 days By comparison With generics . However these differences were outside statistical validity or due to the analysis of small comparison groups, or due to big scatter data V compared ranks. Elongation timing negativity microscopic analyzes on mushrooms at patients With combined damage to the scalp and smooth skin in the treatment with lamiter and terbizil may have been associated with a belated transition to the methodology of N.N. long absence speakers V clinical And microbiological curemycotic process. So, in one of our observations (boy G., 6 years old with a weight body 22 kg) treatment was started lamiter By classical scheme, V connections With no negative results for fungi

he was transferred on the 27th day to treatment with exifin at the same dose, on the 40th day of treatment, the dose was increased to 187.5 mg, and third contract negative analysis on mushrooms was received only on 89th day treatment, i.e. at week 13 of treatment. Probably in this case there was a decrease sensitivity pathogen to drugs terbinafine .

Portability drugs V in general was good. Side effects phenomena V formild diarrheal syndrome occurred in 2 cases (1 at the end of the first week treatment lamiter , 2nd - on second week treatment terbizil ), But They Not demanded that the drug be discontinued.

So the way clinical And mycological permission foci on smooth skin in the treatment of terbinafine and griseofulvin drugs occurred on average behind 3.5 weeks A microsporia hairy parts heads demanded at majority patients at least 5.5 - 7 weeks of systemic and local therapy. Similar terms indicate and others researchers [1-5].

We have calculated the average cost of the drug for the course of treatment of the patient microsporia. In the treatment with griseofulvin, the cost of the drug for a course of treatment patient with microsporia of the scalp amounted to 43 - 66 thousand rubles, and with a combined lesion of the scalp and smooth skin - 45 - 65 thousand rubles; when using lamisil - respectively 122 - 146 thousand rubles and 130 - 156 thousand rubles; when using a lamiter - 48 - 62 thousand and 50 - 60 thousand rubles; at treatment exifin - 54 - 71 thousand And 58 - 70 thousand rubles; A at treatment terbizil – 62 - 77 thousand And 65- 83 thousands rubles.

**Conclusions;** So the way drugs terbinafine may be recommended For treatment microsporia at children And adults on background absence V pharmacy networks griseofulvin . WITH clinical points vision the best results received at use lamisila . WITH pharmacoeconomic points vision most preferred is the treatment of microsporia using lamiter . However, the extension timing stationary treatment children at use generics terbinafine requires further improvement methodologies treatment microsporia, approbation new drugs and methods of external therapy, the development of methods for outpatient treatment these patients.

### **Literature**

1. Anvarovich O. R. et al. ETIOPATHOGENESIS AND CLINICAL COURSE SPECIFIC CHARACTERISTICS OF NEURODERMITIS DISEASE //Web of Scholars: Multidimensional Research Journal. – 2022. – T. 1. – №. 6. – C. 276-280.
2. Bakhodirovich T. I. et al. COMBINED IMMUNOTHERAPY IN THE TREATMENT OF CANDIDIASIAN INFECTIONS OF THE URINARY ORGANS //Web of Scholars: Multidimensional Research Journal. – 2022. – T. 1. – №. 6. – C. 257-263.
3. Baxadirovich T. I. Topical Treatment of Children with Atopic Dermatitis //Texas

- Journal of Medical Science. – 2022. – Т. 8. – С. 132-134.
4. Вахадирович Т. I. AKNE VULGARNIKSNI DAVOLASH SAMARALIGINI BAHOLASH //Eurasian Journal of Medical and Natural Sciences. – 2022. – Т. 2. – №. 3. – С. 41-44.
  5. Аликулова М. Т., Тиллакобилов И. Б. кандиФЛЮ нео при кандидозе крупных складок //Forcipe. – 2020. – Т. 3. – №. 5. – С. 221-222.
  6. Вахадирович Т. I. Dermatological Practice for the Treatment of Adopic Dermatitis and Eczema //Central Asian Journal of Medical and Natural Science. – 2022. – Т. 3. – №. 5. – С. 670-672.
  7. Нарзикулов Р. М. НОВЫЙ ПОДХОД В ЛЕЧЕНИИ ХРОНИЧЕСКОЙ ЭКЗЕМЫ У РАБОТНИКОВ СЕЛЬСКОГО ХОЗЯЙСТВА В УСЛОВИЯХ РЕЗКО КОНТИНЕНТАЛЬНОГО КЛИМАТА (НА ПРИМЕРЕ САМАРКАНДСКОЙ ОБЛАСТИ) //Актуальные аспекты медицинской деятельности. – 2021. – С. 235-240.
  8. Махматкулова Г. М., Тиллакобилов И. Б. СОВРЕМЕННЫЕ МЕТОДЫ ЛЕЧЕНИЯ ТЯЖЕЛЫХ ФОРМ ПСОРИАЗА //Proceedings of International Conference on Modern Science and Scientific Studies. – 2023. – Т. 2. – №. 5. – С. 153-154.
  9. Махматкулова Г. М., Тиллакобилов И. Б. СОВРЕМЕННЫЕ ПРОБЛЕМЫ ДЕРМАТОЗОВ СВЯЗАННЫХ С НАРУШЕНИЕЯМИ ПИТАНИЯ ЗАБОЛЕВАНИЯМИ ЖЕЛУДОЧНО–КИШЕЧНОГО ТРАКТА ПЕЧЕНИ И ЭНДОКРИННЫХ ОРГАНОВ И РАЗРАБОТКА МЕТОДОВ ИХ КОРРЕКЦИИ //Proceedings of International Conference on Modern Science and Scientific Studies. – 2023. – Т. 2. – №. 5. – С. 139-140.
  10. Вахадирович Т. I., Musurmonovich S. M. The Presence of Antibodies to Hiv Infection in Patients First Established Diagnosis of Herpes //Web of Semantic: Universal Journal on Innovative Education. – 2023. – Т. 2. – №. 2. – С. 143-145.
  11. Вахадирович Т. I., Rustamovich R. A. Evaluation of the Efficiency of Sensitivity of N. Gonorrhoeae to Antimicrobial Drugs //Web of Semantic: Universal Journal on Innovative Education. – 2023. – Т. 2. – №. 2. – С. 138-142.
  12. Вахадирович Т. I., Musurmonovich S. M., Nozimovich N. O. Evaluation of Lipid Peroxidation and Enzymes of the Antioxidant System in Patients with Various Forms of Alopecia //Web of Semantic: Universal Journal on Innovative Education. – 2023. – Т. 2. – №. 2. – С. 133-137.
  13. Anvarovich O. R. et al. ETIOPATHOGENESIS AND CLINICAL COURSE SPECIFIC CHARACTERISTICS OF NEURODERMITIS DISEASE //Web of Scholars: Multidimensional Research Journal. – 2022. – Т. 1. – №. 6. – С. 276-280.
  14. Kurbonalievich A. S. et al. UROGENITAL CHLAMYDIOSIS IS A CAUSE OF STERILITY AND CHRONIC PROSTATITIS IN MEN //Web of Scholars: Multidimensional Research Journal. – 2022. – Т. 1. – №. 6. – С. 306-310.
  15. Мансур Т. М., Вохидов Ж. Ж. СОВРЕМЕННОЙ МЕТОДЫ АЛОПЕЦИЯ ДИАГНОСТИКА И ЛЕЧЕНИЕ //SCIENTIFIC APPROACH TO THE MODERN EDUCATION SYSTEM. – 2023. – Т. 2. – №. 14. – С. 225-233.
  16. Мансур Т. М., Вохидов Ж. Ж. СОВРЕМЕННЫЕ МЕТОДЫ ЛЕЧЕНИЯ БОЛЕЗНИ ВИТИЛИГО //SCIENTIFIC APPROACH TO THE MODERN

- EDUCATION SYSTEM. – 2023. – Т. 2. – №. 14. – С. 234-244.
17. Толибов М. М. СОВРЕМЕННОЕ СОСТОЯНИЕ ПРОБЛЕМ ЭПИДЕМИОЛОГИИ, МИКРОБИОЛОГИИ И ТЕРАПИИ АКНЕ (ОБЗОР ЛИТЕРАТУРЫ) //SCIENTIFIC APPROACH TO THE MODERN EDUCATION SYSTEM. – 2023. – Т. 2. – №. 14. – С. 215-224.
  18. Мансур Т. М., Вохидов Ж. Ж. ОПТИМИЗАЦИЯ СОВРЕМЕННЫХ МЕТОДОВ ЛЕЧЕНИЯ ДИФФУЗНОЙ АЛОПЕЦИИ //SCIENTIFIC APPROACH TO THE MODERN EDUCATION SYSTEM. – 2023. – Т. 2. – №. 14. – С. 200-214.
  19. Толибов М. М. и др. ОПТИМИЗАЦИЯ К ЛЕЧЕНИЮ ТЯЖЕЛЫХ ОБОЖЖЕННЫХ БОЛЬНЫХ //SCIENTIFIC APPROACH TO THE MODERN EDUCATION SYSTEM. – 2023. – Т. 2. – №. 14. – С. 190-199.
  20. Abdullaev X. D. et al. EVALUATION OF THE EFFECTIVENESS OF THE DRUG GEPON IN THE TREATMENT OF GENITAL CANDYLOMAS //Web of Scientist: International Scientific Research Journal. – 2021. – Т. 1. – №. 02. – С. 16-19.
  21. Абдуллаев Х. Д. и др. Определение цитокинового статуса у больных с псориазом на фоне стандартного лечения //Общество и инновации. – 2020. – Т. 1. – №. 1/S. – С. 625-629.
  22. Ахмедова М. М., Абдуллаев Х. Д., Камолова М. И. ОЦЕНКА ЭФФЕКТИВНОСТИ МЕТОДОВ ЛЕЧЕНИЯ ОНИХОМИКОЗОВ У ВЗРОСЛЫХ //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – Т. 2. – №. 3. – С. 186-190.
  23. Тошев С. У., Аширов З. Ф., Абдуллаев Х. Д. ЛЕЧЕНИЕ ВИТИЛИГО 308-нм ЭКСИМЕРНЫМ ЛАЗЕРОМ //Актуальные аспекты медицинской деятельности. – 2021. – С. 240-243.
  24. Davlatovich A. X., Haydarjonovna X. S., Raximjon N. S. JYBYK BILAN BO'LGAN PROSTATITNI DAVOLASH USULINI TAKOMILLASHTIRISH //БАРҚАРОРЛИК ВА ЕТАКЧИ ТАДҚИҚОТЛАР ОНЛАЙН ИЛМИЙ ЖУРНАЛИ. – 2022. – С. 359-361.
  25. Ахмедова М., Расулова Н., Абдуллаев Х. Изучение парциальных функций почек у детей раннего возраста с нефропатией обменного генеза //Журнал проблемы биологии и медицины. – 2016. – №. 2 (87). – С. 37-40.
  26. Нарзикулов Р. и др. Принципы терапии у женщин больных гонореей ассоциированные иппп //Журнал вестник врача. – 2019. – Т. 1. – №. 1. – С. 99-102.
  27. Нуруллаева А. А., Рахматова А. Х., Абдуллаев Х. Д. ЗНАЧЕНИЕ МИКРОБНОГО ОБСЕМЕНЕНИЯ КОЖИ ПРИ НЕКОТОРЫХ ЗУДЯЩИХ ДЕРМАТОЗАХ //Молодежь и медицинская наука в XXI веке. – 2019. – С. 125-125.
  28. Abdullaev X. D. et al. EXPERIENCE OF USING THE DRUG GEPON IN THE TREATMENT OF PATIENTS FOCAL SCLERODERMA //ResearchJet Journal of Analysis and Inventions. – 2021. – Т. 2. – №. 03. – С. 60-63.
  29. Абдуллаев Х. Д., Собиров М. С., Жумаева Д. Х. НЕРВНО-ПСИХИЧЕСКИЙ СТАТУС У БОЛЬНЫХ СЕБОРЕЕЙ //Молодежь и медицинская наука в XXI веке. – 2018. – С. 115-116.

30. Ахмедов Ш. К. и др. ЭФФЕКТИВНОСТЬ ЛЕЧЕНИЯ ИЗОТРЕТИНОИНА ПРИ ЛЕЧЕНИИ УГРЕВОЙ БОЛЕЗНИ //Академический журнал Западной Сибири. – 2015. – Т. 11. – №. 1. – С. 56-56.
31. Абдуллаев Х., Толибов М. Allergodermatozlar bilan bog'liq bo'lgan vulgar asneni kompleks davolash Samaraligini o'rganish //Журнал гепатогастроэнтерологических исследований. – 2021. – Т. 2. – №. 3.2. – С. 73-74.
32. Davlatovich A. X., O'gli A. B. X., O'gli I. A. S. BOLALARDA GENITAL GERPESNI DAVOLASH SAMARADORLIGINI VAHOLASH //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 367-369.
33. Абдуллаев Х. Д. и др. ЛАЗЕР В ЛЕЧЕНИИ ВИТИЛИГО //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 495-500.
34. Xolmurodovich D. J., Umidovich N. T., Davlatovich A. X. CLINIC COURSE OF NONSPECIFIC PNEUMONIA //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 510-513.
35. Davlatovich A. X. et al. USING IMMUNOMAX AND 0.1% TACROLIMUS OINTMENT IN THE TREATMENT OF VITILIGO //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 559-562.
36. Davlatovich A. X. et al. EVALUATION OF THE GENERAL SOMATIC STATUS OF PATIENTS WITH VITILIGO BASED ON THE DETERMINATION OF THE CONTENT OF NATURAL ANTIBODIES TO VARIOUS ORGANIS AND TISSUES OF THE BODY //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 472-476.
37. Davlatovich A. X. DETERMINATION OF GENE ACTIVITY IN VITILIGO PATIENTS //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 451-454.
38. Xolmurodovich D. J., Orifovich R. S., Davlatovich A. X. FEATURES OF THE MICROELEMENT STATUS FOR THE DEVELOPMENT OF ATOPIC DERMATITIS IN CHILDREN //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 447-450.
39. Hikmatovich I. N. et al. The use of Fungite in the Local Treatment of Genital Herpes //International Journal of Innovative Analyses and Emerging Technology. – 2021. – Т. 1. – №. 5. – С. 231-232.
40. Davlatovich A. X., Orifovich S. D., Baxtiyorovich A. S. Studying the Relationship of Local Immune Status and the Course of Burn Disease //Central Asian Journal of Medical and Natural Science. – 2022. – Т. 3. – №. 5. – С. 679-682.
41. Davlatovich A. X. VAGINAL TRIXOMONADLAR SHTAMMASINI TRIXOPOLGA VA XIMOTRIPSIN BILAN BILAN SEZGICHLIGINI ANIQLASH //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – С. 645-647.
42. Salamova L. A. et al. Evaluation of the effectiveness of the drug gepon in the treatment of genital candyomas. – 2021.



43. Iskandarovna K. M. SIFILISNING IMMUNOASSAY SHAKLLARI //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – C. 534-536.
44. Iskandarovna K. M. et al. Assessment of the Quality of the Treatment and the Probability of Recurrence of Warts in Children Using Groprenosin //Web of Semantic: Universal Journal on Innovative Education. – 2023. – T. 2. – №. 2. – C. 221-226.
45. Iskandarovna K. M. et al. Quality Evaluation of the Efficiency of the Drug" Roaccutan" in the Treatment of Patients with Acne //Web of Semantic: Universal Journal on Innovative Education. – 2023. – T. 2. – №. 2. – C. 241-245.
46. Iskandarovna K. M., Dilshodovna A. D., Adxamovna A. A. Impact of Socio-Hygienic Conditions of Life on the Health of Students //Central Asian Journal of Medical and Natural Science. – 2022. – T. 3. – №. 5. – C. 644-646.
47. Iskandarovna K. M. et al. Comprehensive Assessment of the Levels of Somatic Health and Disturbances of Adaptation Reserves in Medical Students //Central Asian Journal of Medical and Natural Science. – 2022. – T. 3. – №. 5. – C. 641-643.
48. Iskandarovna K. M., Alamovich K. A., Rabbimovich N. A. Treatment of Urethrogenic Prostatitis Associated with Chlamydia Infection //TA'LIM VA RIVOJLANISH TAHLILI ONLAYN ILMIY JURNALI. – 2021. – T. 1. – №. 5. – C. 44-46.
49. Iskandarovna K. M., Buribaevna I. S., Azamovna A. N. Immunoassay Forms of Syphilis //TA'LIM VA RIVOJLANISH TAHLILI ONLAYN ILMIY JURNALI. – 2021. – T. 1. – №. 5. – C. 47-49.
50. Davlatovich A. X. VAGINAL TRIXOMONADLAR SHTAMMASINI TRIXOPOLGA VA XIMOTRIPSIN BILAN BILAN SEZGICHLIGINI ANIQLASH //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – C. 645-647.
51. Abdullaev X. D. et al. EVALUATION OF THE EFFECTIVENESS OF THE DRUG GEON IN THE TREATMENT OF GENITAL CANDYLOMAS //Web of Scientist: International Scientific Research Journal. – 2021. – T. 1. – №. 02. – C. 16-19.