

RESPIRATORY DYSFUNCTION OF PATIENTS WITH SEVERE BURN INJURY

Usmonov U.R.

Teshayeva D.Ch.

BUKHARA STATE MEDICAL INSTITUTE

ABSTRACT

Dysfunctional changes in the upper respiratory tract in most cases develop as a post-burn pathology. (1,2,12). After a burn, in 30-40% of patients, the general condition worsens and vital organs fail. (3,6,8,11,13,16). The most important factors that damage the mucous membrane of the respiratory tract during IT are various combustion products in a confined space. The development of mucosal edema is primarily dependent on mediators such as histamine, bradykinin, prostacyclin and bronchospasm, and pulmonary vascular resistance is associated with the release of thromboxane and complement components (8.15).

Key words: Multiple organ disorders, burn disease, inhalation injury.

Introduction

Acute respiratory failure in burned patients, regardless of the cause that caused it, often leads to multiple organ failure and mortality. Respiratory failure is one of the most common forms of organ failure in the PON syndrome and, progressing, leads to death in combination with dysfunction of other organs, regardless of the presence of inhalation injury (12,16). The disorder of the respiratory function of the lungs, in turn, disrupts the nutrition of the alveolar tissue, reduces the production of surfactant, contributing to the development of atelectasis (4.14), and increases venous blood shunting (2.5). In burn disease, along with respiratory disorders, non-respiratory functions of the lungs suffer: blood purification from mechanical impurities and their enzymatic destruction, participation in hemostasis and fibrinolysis, water and electrolyte metabolism, metabolism of proteins, fats, biologically active substances (1, 7,9,10).

Purpose of the study: To study the pathology of the respiratory organs in burn disease

Materials and methods of research: For the period 2017-2021, there were 2136 patients in the combustiology department of the Bukhara branch of the RRCEM. Of these, 156 patients had impaired function of the respiratory system. Of these, 96 women (61.5), men 60 (38.5). The age of the patients ranged from 2 to 70 years.

Frank index (IF) more than 60-90 units in 76 (48.7) patients, IF more than 90 units in 80 (51.3%) patients. In 55% burns with flames, 43% with boiling water, in 2% - electrical injury.) To study the dysfunction of the respiratory system in terms of

systemic inflammatory (SIR), we analyzed the clinical indicator of tachypnea and features of the X-ray picture of the lungs (Table No. 1).

Chest radiography is an important diagnostic method for monitoring pulmonary complications and evaluating treatment from day 1 of a burn injury (493).

Results and discussions: First of all, it was noted that among 156 patients with identified disorders of the respiratory system, the vast majority (128 patients) with tachypnea 21-24 in 1 minute had signs of venous plethora of the lungs, expressed to varying degrees and indicating an increase in pressure in the pulmonary artery system. According to the literature data, the pressure in the pulmonary artery is increased during all periods of burn disease, depending on the severity of the burn disease, and the greatest increase in pressure was found in the stages of shock and toxemia with a gradual decrease by the time of healing of burn wounds (11.16).

Hyperserotoninemia plays a role in the genesis of increased pulmonary resistance, hypertension of the pulmonary circulation, impaired perfusion and gas exchange, causing vasoconstriction and bronchial constriction (2,4). The development of venous plethora, and with it tachypnea in burned patients, we associate with manifestations of systemic microcirculation disorders and capillary-alveolar diffusion of blood gases against the background of developed SIRS (17.18.19.20).

Table №.1

Pathological changes	Number of patients B Γ]			groups	Total 156
	1-18	II-40	III - 56	IV-42	
Venous plethora	7	30	52	39	128
Interstitial edema	-	3	10	10	23
Alveolar edema	-	1	3	6	10
ARDSV	-	-	1	2	3
bacterial pneumonia	-	1	8	11	20
Infarct pneumonia	1	5	6	3	15
Atelectasis	1	3	8	6	18
Diaphragm dome position	-	5	7	8	20
Pneumothorax	-	-	1	-	1

Distribution of patients with different severity of burn injury in accordance with x-ray changes in the lungs in most patients, signs of venous plethora appeared on the 1-2 day and persisted for 7-14 days, and in some patients it was much longer during the SVR, which correlated with the severity of the burn injury. With an uncomplicated course of a burn disease, venous plethora, according to radiography, gradually decreased and disappeared, with a complicated one, it appeared again or increased, while some patients developed signs of increased vascular permeability in the

microvessels of the lungs, which manifested itself as interstitial or even alveolar edema. Among 128 patients, more than half (5, 8, 11) had tachypnea combined only with circulatory disorders in the lungs.(22,25)

Signs of increased vascular permeability against the background of venous plethora in the form of interstitial and alveolar pulmonary edema were detected in 23 and 10 patients, respectively. An additional negative role in the formation of edema can be played by excessive infusion therapy.(20)

Conclusions: We assume that interstitial edema, caused by a systemic increase in vascular permeability in SVR, occurred in a larger number of patients with tachypnea, but its severity was minimal and insufficient to detect this sign on radiographs. A more pronounced swelling of the interstitium, and even more so the sweating of edematous fluid into the alveoli, caused more severe dysfunction and even respiratory failure.

Literature:

1. Пулатова, Ш. Х. (2021). АРТЕРИАЛЬНАЯ ГИПЕРТОНИЯ И ХРОНИЧЕСКАЯ СЕРДЕЧНАЯ НЕДОСТАТОЧНОСТЬ: КОМОРБИДНОСТЬ КАК ФАКТОР РИСКА НЕДОСТАТОЧНОЙ ЭФФЕКТИВНОСТИ ТЕРАПИИ. In *АКТУАЛЬНЫЕ ВОПРОСЫ МЕДИЦИНЫ КРИТИЧЕСКИХ СОСТОЯНИЙ* (pp. 59-60).

2. Усмонов, У. Р., & Иргашев, И. Э. (2020). Changes in the morphofunctional properties of thymus and spleen under the influence of mites of different origins. *Новый день в медицине*, (2), 242-244.

3. Влияние вентиляции легких, контролируемой по объему и по давлению, на результаты лечения больных с геморрагическим инсультом / А.И. Грицан, А.А. Газенкамф, Н.Ю. Довбыш, А.В. Данилович // *Вестник анестезиологии и реаниматологии*. — 2012. — № 3. — С.26—31.

4. Rizoyevich, U. U., Olimjonovich, J. O., Khusenovich, S. S., & Sharifboevna, K. D. (2021). Changes in the morphofunctional properties of thymus, spleen and lymphoid system under the influence of mites of different origins. *Web of Scientist: International Scientific Research Journal*, 2(12), 533-540.

5. Пулатова, Ш. Х., Азимов, Б. К., & Тоиров, И. Р. (2019). Эндovasкулярное лечение больных ишемической болезнью сердца. *Евразийский кардиологический журнал*, (S1), 327-328.

6. Rizoyevich, U. U., Olimjonovich, J. O., Khusenovich, S. S., & Sharifboevna, K. D. (2022). CHANGES IN THE MORPHOFUNCTIONAL PROPERTIES OF THYMUS, SPLEEN AND LYMPHOID SYSTEM UNDER THE INFLUENCE OF MITES OF DIFFERENT ORIGINS. *Web of Scientist: International Scientific Research Journal*, 3(1), 23-29.

7. Байханова, М. Б., Бафаев, Ж. Т., & Пулатова, Ш. Х. (2009). Роль врача общей практики в повышении медицинской грамотности населения. *Врач-аспирант*, 28(1), 48-50.

8. Khayotovich, K. D., & Ikromovich, T. I. (2022). SPECIFICITY OF RESUSCITATION MEASURES IN PATIENTS WITH ISCHEMIC HEART DISEASE AND ARRHYTHMIA. *World scientific research journal*, 10(1), 150-155.

9. Хайитов, Д. Х., & Болтаев, Э. Б. (2022). ПОСТРЕАНИМАЦИОН КАСАЛЛИК НАТИЖАСИДА КЕЛИБ ЧИКАДИГАН АСОРАТЛАРНИ БАРТАРАФ ЭТИШДА ЗАМОНАВИЙ ИНТЕНСИВ ТЕРАПИЯ. КЛИНИК АМАЛИЕТДА УЧРАГАН ХОЛАТ. *Academic research in modern science*, 1(9), 172-178.

10. Khayotovich, K. D., & Ikromovich, T. I. (2022). Specific Morpho functional Changes of the Lymphatic System in Patients Suffering from Burns. *Eurasian Research Bulletin*, 15, 81-84.

11. Yarashev A.R., Boltaev E.B., Shabaev Y.K. A retrospective analysis of complications of percutaneous dilated tracheostomy // *New day in medicine*, 2020. 4 (32). P. 301-304.

12. Khayotovich, K.D., & Bekmurodugli, B.E. (2022). Case in clinical practice: Modern intensive care in the treatment of post-resuscitation complications caused by cardiac arrhythmias. *ACADEMICIA: An International Multidisciplinary Research Journal*.

13. Babanazarov, U. T., & Barnoyev, S. S. (2023). Clinical Characteristics of Patients with Chronic Diffuse Liver Disease Against the Background of Covid-19. *Genius Repository*, 26, 49-55.

14. Rizaeva, M. Z. (2022). The clinical course of atrial fibrillation in patients with coronary heart disease. *European journal of molecular medicine*, 2(1).

15. Turobkulovich, B. U., & Khayotovich, K. D. (2024). MORE THAN MINIMUM CONSCIOUSNESS: APPALLIC SYNDROME. *European Journal of Interdisciplinary Research and Development*, 23, 113-115.

16. Ризаева, М. Ж. (2020). ЭФФЕКТИВНОСТЬ И БЕЗОПАСНОСТЬ ЭЛЕКТРИЧЕСКОЙ КАРДИОВЕРСИИ ПРИ ПЕРСИСТИРУЮЩЕЙ ФОРМЕ ФИБРИЛЛЯЦИИ ПРЕДСЕРДИЙ. *Новый день в медицине*, (4), 322-325.17. Потапов А.А., Крылов В.В., Лихтерман Л.Б. и др. Современные рекомендации по диагностике и лечению тяжелой черепно-мозговой травмы // *Журнал вопросы нейрохирургии*. – 2006. – № 1. – С. 3–8.

18. Qoyirov, A. Q., Kenjaev, S. R., & Xaitov, S. S. (2020). Egamova NT, Boltaev EB The role of delirium in patients with myocardial infarction of complicated acute heart failure. *New Day in Medicine*, 3(31), 68-71.

19. Kh, P. S., & Ganiev, N. S. (2022). The Importance of Cardioprotective Artificial Ventilation of The Lungs in Intensive Care. *Eurasian Research Bulletin*, 15, 208-212.

20. Эшонов, О. Ш., & Болтаев, Э. Б. (2020). СПОСОБ ЭКСТРЕННОГО ОПРЕДЕЛЕНИЯ СТЕПЕНИ ТЯЖЕСТИ ЭНДОТОКСИКОЗА ПРИ НЕОТЛОЖНЫХ СОСТОЯНИЯХ. *Новый день в медицине*, (1), 462-464.

21 Sharifovich, B. S., & Hayotovich, X. D. (2023). Management Of Deep Vein Thrombosis. *Genius Repository*, 27, 59-71.

22. Ураков, Ш. Т., & Ризаева, М. Ж. (2019). КЛИНИЧЕСКИЙ СЛУЧАЙ ПАЦИЕНТА С СИНДРОМОМ МАРФАНА. *Новый день в медицине*, (4), 439-440.

23. Мирзажоннова, Г. С., Пулатова, Ш. Б., & Набиева, Д. А. (2023). *Частота поражения сердца при анкилозирующем спондилите* (Doctoral dissertation, Zamonaviy tibbiyotning dolzarb muammolari yosh olimlar xalqaro anjumani, Uzbekiston).

24. Бабаназаров, У. Т., & Хайитов, Д. Х. (2024). БОЛЬШЕ, ЧЕМ МИНИМАЛЬНОЕ СОЗНАНИЕ: АПАЛЛИЧЕСКИЙ СИНДРОМ. *European Journal of Interdisciplinary Research and Development*, 23, 109-112.

25. Sh, B. S. (2024). IMPORTANCE OF MYORELAXATION IN PATIENTS WITH SEVERE COMA. *Journal of new century innovations*, 47(3), 5-8.

26. Babanazarov, U. T., & Barnoyev, S. S. (2023). Clinical Characteristics of Patients with Chronic Diffuse Liver Disease Against the Background of Covid-19. *Genius Repository*, 26, 49-55.